

**FUZZY FUNCTIONAL ANALYTIC STUDY OF STATISTICAL CONVERGENCE
IN VIEW OF FUZZY SEQUENCE SPACE**

ANINDITA BASU

Abstract. The aim of this work is to extend and study the idea of fuzzy statistical convergence, fuzzy lacunary convergence and fuzzy lacunary statistical convergence for the fuzzy sequence space $F(L(\mathbb{R}), M, p, s)$ of the real valued fuzzy number sequences by utilizing the modern techniques of fuzzy functional analysis. This sequence space is an unified approach of the class of sequences of fuzzy numbers $L(\mathbb{R})$, where an Orlicz function M , a normal sequence space F , a bounded sequence of strictly +ve real numbers $p = \{p_k\}$ with $\inf p_k \geq 0$ and a multiplier sequence $(k^{-s}), s \geq 0$ have been used. Further, some important results regarding these type of convergences and their inter-relations have been attempted to derive which are analogous to the results provided by the classical analysis.

Keywords: fuzzy statistical convergence, fuzzy lacunary convergent space, fuzzy lacunary statistical convergence, para-normed space, normal space.

AMS subject classification: 40A05, 40A35, 40C05, 40D05, 46A45.

Introduction: Fuzzy number sequence space is one of the active areas of research of fuzzy functional analysis. The concept of fuzzy sets and fuzzy set operations were first introduced by Zadeh [17] and Matloka [18] first introduced bounded and convergent sequences of fuzzy numbers and studied their properties. Later on, applying the notion of fuzzy real numbers, fuzzy real-valued sequence space was first introduced and studied by Nanda [21]. He showed that the set of all convergent sequences of fuzzy numbers forms a complete metric space. Later on the study of theory of fuzzy sequence spaces has been continued by Nanda and Tripathy [22], Savas [9], Mursaleen et al. [19], Fang et al. [16], Talo et al. [23, 24], Dutta et al. [13], Esi et al. [28], Tripathy et al [7, 8], Basu et al [1] and many others.

Nuray and Savas [11] first introduced the notion of statistically convergent and statistically Cauchy sequence for sequence of fuzzy numbers. Later on Nurray [12] studied a related concept, namely, lacunary convergent and lacunary statistical convergent sequences of Fuzzy numbers and various inclusion relations. Recently the notion of Statistically convergent, lacunary statistical convergent sequences of fuzzy numbers defined by modulus and Orlicz function have been studied by Savas ([10]), Dutta et al. [13], Esi et al [4], Sarma [5], Tripathy et al [6] and many authors. Later on Mursaleen and Basarir [19], Fang and Huang [29] and many other authors studied the different convergence methods for fuzzy single and double sequences.

A sequence of positive integers $\theta = (k_r)$ is called 'lacunary' if $k_0 = 0, 0 < k_r < k_{r+1}$ and $h_r = k_r - k_{r-1} \rightarrow \infty$ as $r \rightarrow \infty$. The intervals determined by θ are denoted by I_r , where $I_r = (k_{r-1}, k_r]$ and the ratio $\frac{k_r}{k_{r-1}}$ is denoted by q_r .

The space of lacunary strong convergence N_θ has been introduced by Freedman et al. [3]. Later on Nurray [16] studied a related concept, namely, lacunary convergent and lacunary statistical convergent sequences of Fuzzy numbers and various inclusion relations.

The present author has made an attempt to fuzzify the concepts of statistical and lacunary statistical convergence methods on the generalized composite class $F(L(\mathbb{R}), M, p, s)$ of fuzzy Orlicz sequence spaces introduced by Basu et al. [2].

The class $F(L(\mathbb{R}), M, p, s)$: Let (F, g_F) be a normal sequence space with paranorm g_F which satisfies the following properties:

- i) g_F is monotone and;
- ii) co-ordinatewise convergence implies convergence in paranorm g_F i.e., if $(X^n) = (X_k^n) \in F$, then $X_k^n \rightarrow 0$ as $n \rightarrow \infty \implies g_F(X^n) \rightarrow 0$ as $n \rightarrow \infty$

Let M be a Orlicz function. We define

$$F(L(\mathbb{R}), M, p, s) = \left\{ X = (X_k) : X_k \in L(\mathbb{R}), \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \in F \text{ for some } \rho > 0 \right\}$$

where $\{p_k\}$ is a bounded sequence of strictly +ve real numbers with $\inf p_k > 0$ and $s \geq 0$.

We define a metric topology on this fuzzy sequence space $F(L(\mathbb{R}), M, p, s)$ as follows:

For $X = (X_k), Y = (Y_k) \in F(L(\mathbb{R}), M, p, s)$,

$$d(X, Y) = \inf \left\{ \rho^{p_k/T} > 0 : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k, Y_k)}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \right\}$$

where $T = \max(1, H), H = \sup_k p_k < \infty, \inf_k p_k > 0$.

Particular Cases : It is shown that this generalized class of sequences of fuzzy real numbers gives rise to many well known fuzzy sequence spaces on specifying the sequence space F , the Orlicz function M , the bounded sequence $\{p_k\}$ of positive real numbers and $s \geq 0$ as follows:

1. If we take $F = c, \ell_p, s = 1, p_k = 1$, for each $k \in \mathbb{N}, M(x) = x$, for all $x \in [0, \infty)$, $F(L(\mathbb{R}), M, p, s)$ reduces to the spaces of Nanda [21];
2. If we take $F = w(p), c_0(p), s = 1, M(x) = x$, for all $x \in [0, \infty)$, $F(L(\mathbb{R}), M, p, s)$ reduces to the spaces of Mursaleen et al. [22];
3. If we take $F = \ell(p)$ [26], $s = 1, M(x) = x$, for all $x \in [0, \infty)$, $F(L(\mathbb{R}), M, p, s)$ reduces to the spaces of Nurray and Savas [16];
4. If we take $F = \ell_\infty, c, c_0, \ell_p, M(x) = x$, for all $x \in [0, \infty), s = 1, p_k = 1$, for each $k \in \mathbb{N}$, $F(L(\mathbb{R}), M, p, s)$ gives rise to the space of Talo et al. [23].

The following definitions and preliminary concepts have been used throughout the work

Definitions and Propositions:

Definition 1. (Paranorm)[15] Let X be a linear space. Then $g: X \rightarrow \mathbb{R}$ is called a paranorm on X if for $x, y \in X$ and any scalar λ , (i) $g(x) \geq 0$; (ii) $x = \theta \Rightarrow g(x) = 0$; (iii) $g(x) = g(-x)$; (iv) $g(x + y) \leq g(x) + g(y)$; (v) $g(\lambda_n x_n - \lambda x) \rightarrow 0$ as $n \rightarrow \infty$ whenever $\lambda_n \rightarrow \lambda$ and $x_n \rightarrow x$ for scalars λ_n, λ and $x_n, x \in X$.

Definition 2. (Monotone paranorm)[15] A paranorm g on a scalar valued paranormed sequence space Z is called monotone paranorm if $X = (x_k) \in Z, Y = (y_k) \in Z$ and $|x_k| \leq |y_k|$ implies $g(x) \leq g(y)$. The space (Z, g) is called monotone paranormed space.

Definition 3.[22] A sequence space λ of sequences fuzzy numbers is said to be solid (or normal) if $(Y_n) \in \lambda$, whenever $\bar{d}(Y_n, \bar{0}) \leq \bar{d}(X_n, \bar{0})$ for some $(X_n) \in \lambda$.

Definition 4. [22] A metric \bar{d} is said to be translation invariant metric on $L(\mathbb{R})$ if

$$\bar{d}(X + Z, Y + Z) = \bar{d}(X, Y) \text{ for } X, Y, Z \in L(\mathbb{R}).$$

Proposition 1. [22] Let $X, Y, Z, W \in L(\mathbb{R})$ and $k \in \mathbb{N}$. Then,

- (i) $(L(\mathbb{R}), \bar{d})$ is a complete metric space [20];
- (ii) $\bar{d}(kX, kY) = |k| \bar{d}(X, Y)$
- (iii) $\bar{d}(X + Y, W + Z) \leq \bar{d}(X, W) + \bar{d}(Y, Z)$;

Proposition 2. [22] Let $X, Y, Z, W \in L(\mathbb{R})$ and $k \in \mathbb{N}$. Then

- (i) If $X_k \rightarrow 0$ as $k \rightarrow \infty$ then $\bar{d}((X_k, \bar{0})) \rightarrow \bar{d}((X, \bar{0}))$ as $k \rightarrow \infty$;

Proposition 3. [16] Let (p_k) be a bounded sequence of strictly positive real numbers with

$0 < p_k \leq \sup_k p_k = H, D = \max(1, 2^{H-1}), T = \max(1, H)$. Then

1. $|a_k + b_k|^{p_k} \leq D\{|a_k|^{p_k} + |b_k|^{p_k}\}$;
2. $|\lambda|^{p_k} \leq \max(1, |\lambda|^H)$;

Natural Density of a set: The natural density of a set A is defined by

$$\delta(A) = \lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n: k \in A\}|, \text{ where}$$

the vertical bar denotes the cardinality of the set enclosed.

Orlicz Function: [20]: An Orlicz function is a function $M: [0, \infty) \rightarrow [0, \infty)$ which is continuous, non-decreasing and convex with $M(0) = 0, M(x) > 0$ for $x > 0$, and $M(x) \rightarrow \infty$ as $x \rightarrow \infty$.

An Orlicz function M can always be represented in the integral form $M(x) = \int_0^x q(t) dt$, where q , known as the Kernel of M , is right differentiable for $t \geq 0, q(0) = 0, q(t) > 0$ for $t > 0, q(t) \rightarrow \infty$ as $t \rightarrow \infty$.

An Orlicz function is said to satisfy Δ_2 -condition for all values of x , if there exists a constant $K > 0$, such that $M(2x) \leq K M(x)$ for all $x \geq 0$. This condition is equivalent to $M(Lx) \leq KL M(x)$ for all values of $x \geq 0$ and for all $L \geq 1$ and $M(\lambda x) \leq \lambda M(x)$ for all λ with $0 < \lambda < 1$.

Fuzzy Number: [18]:

A fuzzy number X is a mapping $X: \mathbb{R} \rightarrow [0, 1]$ satisfying

- (i) X is normal, i.e., \exists an $t_0 \in \mathbb{R}$ such that $X(t_0) = 1$;
- (ii) X is convex, i.e., $X(t) \geq X(s) \wedge X(r) = \min(X(s), X(r))$, where $r < t < s$;
- (iii) X is upper semicontinuous i.e., for each $\epsilon > 0, X^{-1}([0, a + \epsilon])$ is open in the usual topology of \mathbb{R} , for all $a \in [0, 1]$;
- (iv) $X^0 = \{t \in \mathbb{R}: X(t) > 0\}$ is compact.

In the present work, being motivated by the existing literature of statistical convergence of sequences of fuzzy numbers, the present author has defined the statistical convergence and lacunary statistical convergence on $F(L(\mathbb{R}), M, p, s)$ as follows:

Statistical Convergence on $F(L(\mathbb{R}), M, p, s)$: A fuzzy number sequence $X = (X^{(n)}) = ((X_k^{(n)}))$ in $F(L(\mathbb{R}), M, p, s)$ is said to converge statistically to $X_k = (X_1, X_2, \dots)$, $X_k \in L(\mathbb{R})$, if for any $\epsilon > 0$,

$$\delta \left(\left\{ n \in \mathbb{N} : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^{(n)} - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \geq \epsilon \right\} \right) = 0$$

for each $k \in \mathbb{N}$. It is denoted by $x^{(n)} \xrightarrow{stat} L$.

A fuzzy number sequence $X = (X^{(n)}) = ((X_k^{(n)}))$ in $F(L(\mathbb{R}), M, p, s)$ is said to be statistically Cauchy if for any $\epsilon > 0, \exists$ a natural number $m = m(\epsilon)$ such that

$$\delta \left(\left\{ n \in \mathbb{N} : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \geq \varepsilon \right\} \right) = 0 \text{ for } n \geq m \text{ and for each } k \in \mathbb{N}.$$

Lacunary Convergence and Lacunary Statistical Convergence on $F(L(\mathbb{R}), M, p, s)$

The fuzzy number sequence $X = (X^{(n)}) = ((X_k^{(n)}))$ in $F(L(\mathbb{R}), M, p, s)$ is said to be $N_\theta(L(\mathbb{R}), M, p, s)$ convergent to $X = (X_k)$ if for every $\varepsilon > 0$,

$$\lim_{r \rightarrow \infty} \frac{1}{h_r} \sum_{k \in I_r} \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \geq \varepsilon = 0 \text{ for } n=1, 2, \dots$$

The fuzzy number sequence $X = (X^{(n)}) = ((X_k^{(n)}))$ in $F(L(\mathbb{R}), M, p, s)$ is said to be $S_\theta(L(\mathbb{R}), M, p, s)$ -convergent to $X = (X_k)$ if for every $\varepsilon > 0$,

$$\lim_{r \rightarrow \infty} \frac{1}{h_r} C \left(\left\{ n \in \mathbb{N} : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \geq \varepsilon \right\} \right) = 0 \text{ for } n=1, 2, \dots$$

Results

Theorem. If $X = (X^{(n)}) = ((X_k^{(n)}))$ be a fuzzy number sequence in $F(L(\mathbb{R}), M, p, s)$, then the following conditions are equivalent:

- (a) $X^{(i)} \xrightarrow{stat} X$;
- (b) There exists $Y^{(i)} = ((Y_k^{(i)}))$ and $Z^{(i)} = ((Z_k^{(i)}))$ in $F(L(\mathbb{R}), M, p, s)$ such that $X^{(i)} = Y^{(i)} + Z^{(i)}$ and $\lim_{i \rightarrow \infty} Y^{(i)} = X$ in $L(\mathbb{R})$ and $Z^{(i)} \xrightarrow{stat} \bar{0}$;
- (c) There is a subsequence $I = \{i_r\}$ of \mathbb{N} such that $\delta(I) = 1$ and $\lim_{r \rightarrow \infty} X^{(i_r)} = X$ in $L(\mathbb{R})$ where $X = (X_1, X_2, \dots) \in L(\mathbb{R})$.

Proof: (a) \Rightarrow (b):

Let us select an increasing sequence of positive integers, where $N_0 = 0, N_0 < N_1 < N_2 < \dots$ and if $i > N_j$,

$$\delta \left(\left\{ i \in \mathbb{N} : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^i - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \geq \frac{1}{j} \right\} \right) < 1/j \text{ for each } k \in \mathbb{N}.$$

We define $Y^{(i)}$ and $Z^{(i)}$ as follows:

If $N_0 < k < N_1, Z_k^{(i)} = \bar{\theta}$ and $Y_k^{(i)} = X_k^{(i)}$ for each k .

If $N_m < i < N_{m+1}, m \geq 1$, set

$$Y_k^{(i)} = X_k^{(i)}, Z_k^{(i)} = \bar{\theta} \quad \text{if } \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} < \frac{1}{m} \text{ and}$$

$$Y_k^{(i)} = X_k, Z_k^{(i)} = X_k^{(i)} - X_k \text{ if } \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \geq \frac{1}{m}$$

Then $X_k^{(i)} = Y_k^{(i)} + Z_k^{(i)}$ for each i and k .

Claim: $\lim_{i \rightarrow \infty} Y_k^{(i)} = X_k$.

It follows from: For arbitrary $\varepsilon > 0$ we choose m so that $\varepsilon > \frac{1}{m}$.

Then for $i > N_m$, as $X^{(i)} \xrightarrow{stat} X$ holds, by construction

$$\left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} < \frac{1}{m}$$

Since ε is arbitrary, the claim is justified.

Claim: $Z^{(i)} \xrightarrow{stat} \bar{0}$.

Let $\eta > 0$ and $m \in \mathbb{N}$ such that $\frac{1}{m} < \eta$.

It is trivially true that

$$\delta \left(\left\{ i \in \mathbb{N} : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(Z_k^i, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \neq 0 \right\} \right) \geq \delta \left(\left\{ i : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(Z_k^i, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \geq \varepsilon \right\} \right)$$

Let $\eta > 0$ and $m \in \mathbb{N}$ such that $\frac{1}{m} < \eta$.

Then from the construction it follows that, if $N_m < i \leq N_{m+1}$, then $Z_k^i \neq \bar{0}$ if and only if

$$\left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^i - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} > \frac{1}{m}$$

So, if $N_p < i < N_{p+1}$, $p \geq 1$,

$$\left\{ i \in \mathbb{N} : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(Z_k^i, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} \neq 0 \right\} \subseteq \left\{ i \in \mathbb{N} : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^i - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} > \frac{1}{p} \right\}$$

Consequently, if

$N_p < i < N_{p+1}$, $p > m$, since M is non-decreasing, it follows that

$$\delta \left(\left\{ i \in \mathbb{N} : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(Z_k^i, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} \neq 0 \right\} \right)$$

$$\leq \delta \left(\left\{ i \in \mathbb{N} : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^i - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} > \frac{1}{p} \right\} \right) < \frac{1}{p} < \frac{1}{m} < \eta$$

And our claim is justified.

Proof: (b) \Rightarrow (c) : Let (b) hold and let us define $I = (i_r)$ to be a subsequence of \mathbb{N} such that $i \in I$ if and only if for any $\varepsilon > 0$,

$$\delta \left(\left\{ i \in \mathbb{N} : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(Z_k^i, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} \geq \varepsilon \right\} \right) = 0$$

$$\text{Then } \delta(I) = \delta \left(\mathbb{N} - \left\{ i \in \mathbb{N} : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(Z_k^i, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} < \varepsilon \right\} \right) = 1 - 0 = 1.$$

Since $X_k^{(i_r)} = Y_k^{(i_r)}$ and $\lim_{r \rightarrow \infty} Y_k^{(i_r)} = X_k$, by the given condition, (c) holds.

Proof: (c) \Rightarrow (a) : Let (c) hold .

Then for arbitrary $\varepsilon > 0$, there exist a subsequence $I = \{i_r\}$ of \mathbb{N} such that $\delta(I) = 1$ and $m = m(\varepsilon) \in \mathbb{N}$ such that

$$\delta \left(\left\{ i \in \mathbb{N} : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^{(i_r)} - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} < \varepsilon \right\} \right) \text{ for } r \geq m.$$

Then the result follows from the fact that

$$\delta \left(\left\{ i \in \mathbb{N} : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^i - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{\frac{1}{T}} \geq \varepsilon \right\} \right)$$

$$< \delta(\mathbb{N} - \{i_r : r > m\})$$

$$= 1 - \delta(\{i_r : r > m\})$$

$$= 1 - 1$$

$$= 0$$

Theorem. Let $\theta = (k_r)$ be a lacunary sequence. Then for $((X_k^n)) \in N_\theta(L(\mathbb{R}), M, p, s)$ and $(X_k) \in L(\mathbb{R})$,

(i) $X_k^n \rightarrow X_k$ in $\in N_\theta(L(\mathbb{R}), M, p, s) \Rightarrow X_k^n \rightarrow X_k$ in $\in S_\theta(L(\mathbb{R}), M, p, s)$

(ii) $((X_k^n)) \in \ell_\infty(L(\mathbb{R}), M, p, s)$ and if $X_k^n \rightarrow X_k$ in $\in S_\theta(L(\mathbb{R}), M, p, s)$ then $X_k^n \rightarrow X_k$ in $\in N_\theta(L(\mathbb{R}), M, p, s)$.

(iii) $\ell_\infty(L(\mathbb{R}), M, p, s) \cap N_\theta(L(\mathbb{R}), M, p, s) = \ell_\infty(L(\mathbb{R}), M, p, s) \cap S_\theta(L(\mathbb{R}), M, p, s)$

Proof: (i) \Rightarrow (ii) : The result holds from the following inequalities

$$\begin{aligned} \frac{1}{h_r} \sum_{k \in I_r} \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} &\geq \frac{1}{h_r} \sum_{k \in I_r} [\varepsilon]^{1/T} \\ &\geq \frac{1}{h_r} \sum_{k \in I_r} \min[\varepsilon^h, \varepsilon^H]^{1/T} \\ &\geq \frac{1}{h_r} \sum_{k \in I_r} \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \geq \varepsilon \\ &\geq \frac{1}{h_r} \min[\varepsilon^h, \varepsilon^H]^{1/T} \left\{ k \in I_r : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \right\} \end{aligned}$$

for $0 < h = \inf p_k \leq p_k \leq H = \sup p_k$.

Proof: (ii) ⇒ (iii): The result holds from the following inequalities $\frac{1}{h_r} \sum_{k \in I_r} \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T}$

$$\begin{aligned} &= \frac{1}{h_r} \sum_{k \in I_r} \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \\ &\quad \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \geq \varepsilon \\ &\quad + \frac{1}{h_r} \sum_{k \in I_r} \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \\ &\quad \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} < \varepsilon \\ &\leq \frac{1}{h_r} \sum_{k \in I_r} \max[K^h, K^H]^{1/T} + \frac{1}{h_r} \sum_{k \in I_r} [\varepsilon]^{1/T} \\ &\quad \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \geq \varepsilon \quad \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} < \varepsilon \\ &\leq \max[K^h, K^H]^{1/T} \frac{1}{h_r} \left\{ k \in I_r : \left[g_F \left(k^{-s} \left[M \left(\frac{\bar{d}(X_k^n - X_k, \bar{0})}{\rho} \right) \right]^{p_k} \right) \right]^{1/T} \geq \varepsilon \right\} + [\max(\varepsilon^h, \varepsilon^H)]^{1/T} \end{aligned}$$

for $0 < h = \inf p_k \leq p_k \leq H = \sup p_k$.

Proof: (iii) ⇒ (i): Follows from (i) ⇒ (ii) and (ii) ⇒ (iii)

References:

1. Basu, "Fuzzy composite Modular sequence space", Mathemaical Sciences International Research Journal, Vol-2, Issue-2, 2013,229-235.
2. Basu, & H. Dutta, "Some Fuzzy Sequence Spaces defined by Orlicz functions", under revision.
3. R. Freedman, , J. J.,Sember, M. Raphel, Some Cesaro type Summability, Proc. Lon. Math. Soc., (3), 37. 1978, 508-520.
4. Esi., M. Acikgoz, "Some classes of difference sequences of fuzzy numbers defined by a sequence of moduli",Acta mathematica Scientia vol. 31 (B), 2011, pp. 229–236.
5. Sarma, "On a class of sequences of fuzzy numbers defined by modulus function", Int. J. Sci. Tech. vol.2,2007, pp. 25–28.
6. Dr.K.V.Radha, Mass Transfer Correlation for Decolorization Studies in Immobilized Packed Bed Bioreactor; Engineering Sciences international Research Journal: ISSN 2320-4338 Volume 3 Issue 1 (2015), Pg 17-19
7. C. Tripathy, and B.Sarma, "Some double sequence spaces of fuzzy numbers defined by Orlicz functions", Acta Math. Scient. vol. 31(B), 2011, pp. 134–140..
8. C.Tripathy, and A. J. Dutta, "On Fuzzy real-valued double sequence spaces", Sochow J. Maths. vol.32,2006, pp. 509–520.
9. C.Tripathy, and A. J. Dutta, "On fuzzy real-valued double sequence space ${}_2\ell_F^p$.", Math. Comput. Model vol. 46, 2007, pp. 1294–1299.
10. Savas, "A note on sequence of fuzzy numbers", Inform Sci. vol. 124, 2000, pp. 297–300.

11. Savas, strongly summable sequences spaces in 2-normed space defined by ideal convergence and an
12. Orlicz function, *Appl. Math. and Comp.*, 217, (2010), 271-276
13. M. Venkateswarlu , B. Rajini Kanth, A Simple Method for the Determination of Magneto Rheological Properties of Nanoparticulated Cobalt Ferrite Based Magnetorheological Fluids; *Engineering Sciences international Research Journal: ISSN 2320-4338 Volume 3 Issue 1 (2015)*, Pg 20-23
14. Nurray, and E. Savas, "Statistical convergence of sequences of fuzzy numbers", *Mathematica Slovaca* vol. 45, 1995, pp. 269-273
15. Nurray, Lacunary Statistical Convergence of sequences of fuzzy numbers, *Fuy sets and systems*, 99 (1998), 353-356.
16. Dutta., A characterization of the class of statistically pre-Cauchy double sequences of fuzzy numbers, *Applied Mathematics and Information Sciences*, 7 (4) (2013), pp. 1437-1440.
17. J. Maddox, "Sequence spaces defined by a modulus", *Proc. Camb. Phil. Soc.* vol. 100, 1986, pp. 161-166.
18. Arnab Acharyya, Application of Image Halftone Technique in Visual Secret Sharing of Text Image; *Engineering Sciences international Research Journal: ISSN 2320-4338 Volume 3 Issue 1 (2015)*, Pg 24-27
19. J. Maddox, "Elements of Functional Analysis", Cambridge University press, Cambridge, (1970).
20. J. Fang, H. Huang, "On the level convergence of a sequence of fuzzy numbers", *Fuzzy Sets Sys.* vol. 147, 2004, pp. 417-435.
21. A. Zadeh, "Fuzzy sets", *Inform. Control* vol. 8, 1965, pp. 338-353.
22. Ar. Arpan Dasgupta, Dr. Madhumita Roy, Comparing Visual Comfort in Daylight Pattern Between An Old and A Modern office Building; *Engineering Sciences international Research Journal: ISSN 2320-4338 Volume 3 Issue 1 (2015)*, Pg 28-35
23. Matloka, "Sequences of fuzzy numbers, *BUSEFAL* vol. 28, 1986, pp. 28-37.
24. Mursaleen, and M. Basarir, On some new sequence spaces of fuzzy numbers", *Indian J. Pure Appl. Math.* vol. 34, 2003, pp. 1351-1357.
25. P. K. Kamthan, and M. Gupta, "Sequence spaces and Series", Marcel Dekker, New York, 1981.
26. S. Nanda, "On sequences of fuzzy numbers", *Fuzzy Sets Sys.* vol. 33, 1989, pp. 123-126.
27. O. Talo, and F. Basar, "Determination of the duals of classical sets of sequences of fuzzy numbers and related matrix transformations", *Comput. Math. Appl.* vol. 58, 2009, pp. 717-733.
28. S. Nanda, and B. K. Tripathy, "Absolute value of fuzzy real numbers and fuzzy sequence spaces", *J. Fuzzy Math.* vol. 8, 2000, pp. 883-892.
29. Talo, and F. Basar, "Certain spaces of sequences of fuzzy numbers defined by a modulus function", *Demonstratio Mathematica* vol. XLIII, 2010, pp. 139-149.
30. Dhaneswar Kalita, Sub-Linear Dc Photoconductivity of thermally Evaporated Polycrystalline CDSE Thin Films; *Engineering Sciences international Research Journal: ISSN 2320-4338 Volume 3 Issue 1 (2015)*, Pg 36-44

Anindita Basu

Assistant Professor, Department of Mathematics,

Dr. B. N. Dutta Smriti Mahavidyalaya, Burdwan-713407, West Bengal, India.