

## SUGARCANE SEEDLING ROOTING MEDIA AND SEEDLING VIGOUR

CH. MUKUNDA RAO, M. BHARATHALAKSHMI, K. VEERABHADRA RAO,  
N. VENUGOPALA RAO

**Abstract:** A ploy tray experiment was conducted with different rooting media and raised sugarcane seedlings at RARS, Anakapalle with RBD of four replications during sugarcane cropping season of 2014-15 and 2015-16. The results of the experiment revealed that the seedling vigour index (SVI) of sugarcane was better with the sugarcane seedlings raised in rooting medias of coir pith + vermi compost (1:1) (T<sub>1</sub>) + Urea 5 kg per 100 kg media (SVI 37.95), T<sub>1</sub> + Sett dipping with 10% lime solution (SVI 33.36) and T<sub>1</sub> + Sett treatment with GA<sub>3</sub> 100 PPM (SVI 30.46). The seedlings performed better with high seedling vigour index which can sustain well under transplanted conditions / fields. The additional nourishment through Nitrogen in Urea, Calcium content of 10% lime solution and plant growth substance (GA<sub>3</sub>) helped in the respective treatments for better seedling vigour index over other treatments studied. Sugarcane rooting media (vermin compost + coir pith) is rich in Nitrogen (0.864%), Phosphorous (0.82%), Potassium (1.16%), Iron (4,220 ppm) and Manganese (1200 ppm).

**Keywords:** Sugarcane single node seedling, polytray, rooting media, GA<sub>3</sub>, IAA, Seedling vigour index (SVI), Coir pith and vermin compost.

**Introduction:** Sugarcane is normally propagated by stock cuttings consisting of 2-3 bud setts. In commercial system of sugarcane planting nearly 6-8 tonnes of seed cane per hectare is used as planting material. Establishing the sugarcane crop using bud chip / single node in place of setts could save about 80% by weight of stock material (Radha Jain and Solomon, 2010 a). Indian sugarcane experts Narsimha Rao (1977) and Ramaiah et al.,(1977) showed the feasibility of eliminating the internode part of the seed piece and using only buds for commercial planting. Sugarcane seedling transplanting expected to reduce overall pressure on water resources and contributing to recovering of eco system in long run. Seedling transplanting is an alternative to conventional seed - water and space intensive sugarcane cultivation. Seedling transplanting in sugarcane (Single node technology) is becoming very popular and sugar industry is also encouraging due to its advantage on reduced cost of cultivation and higher yields. Moreover the seed cane : out put ratio is improved to 1:40 compared to that of 1:10 in conventional planting (Prasada Rao,2015) and it gives better yield in ratoon crop (Solomon,2015). In addition Naidu et al., 2015 reported that use of single bud setts also reduces seed cane rate and there by reduces cost of cultivation. But there is no information on standard rooting media for raising cane seedlings. Different people are using as their own knowledge for rooting media preparation thinking rooting media as a support medium for raising sugarcane seedlings. Some people are using enriched nutrient medium and some are using with out any nutrient enrichment. So, it is essential to take up study on standardization of rooting media for raising sugarcane seedlings under a scientific manner. Under this background a trial was conducted for two

years at Regional Agricultural Research Station, Anakapalle.

**Materials and Methods:** A poly tray experiments was conducted at Regional Agricultural Research Station, Anakapalle in the net house during 2014-15 and 2015-16 sugarcane cropping season with date of planting in January month. The experiment was conducted under RBD design with four replications (each poly tray as one replication) duly following procedure of raising single node seedling in Principal Scientist (Crop Physiology), RARS, Anakapalle - 531 001, A.P.

sugarcane. The following are the treatments under the poly tray experiment. T<sub>1</sub>- Coir pith + Vermi compost (1:1); T<sub>2</sub>- T<sub>1</sub> + DAP (5kg / 100 kg media); T<sub>3</sub> - **T<sub>1</sub> + Urea (5kg / 100 kg media)**; T<sub>4</sub>- T<sub>1</sub> + DAP (2.5 kg / 100 kg media) + Urea (2.5 kg / 100 kg media); T<sub>5</sub>- T<sub>1</sub> + Sett treatment with CaCl<sub>2</sub> (2g/lit) + Ethrel (2ml/lit); T<sub>6</sub>- T<sub>1</sub> + ZnSo<sub>4</sub> (0.5 kg) + FeSo<sub>4</sub> (1.0 kg) /100 kg media ; T<sub>7</sub>- T<sub>1</sub> + Sett treatment CaCl<sub>2</sub> (2g/lit) + Urea (10g/lit); T<sub>8</sub>- T<sub>1</sub> + Ethephan (2ml/lit); T<sub>9</sub>- **T<sub>1</sub> + Sett treatment with 10 % lime solution**; T<sub>10</sub>- **T<sub>1</sub> + Sett treatment with 100 ppm GA<sub>3</sub>**; T<sub>11</sub>- T<sub>1</sub> + Sett treatment with IAA 50 ppm; T<sub>12</sub>- T<sub>1</sub> + Sett treatment with DM 45 + Chlorpyriphos; T<sub>13</sub>- Progressive farmers rooting media.

The seedlings were grown, observed upto 45 days and data was taken duly following standard procedures. The seedling vigour index was calculated based on the formulae seedling length multiplied by per cent germination and divided by 100. Data was statistically analyzed by following standard statistical methods. (Panse and Sukhatme, 1978)

**Results and Discussions:** Sugarcane rooting media (vermin compost + coir pith) is rich in Nitrogen (0.864%), Phosphorous (0.82%), Potassium (1.16%), Iron (4,220 ppm) and Manganese(1200 ppm)(Table 1).

**Table1: Nutrient composition of rooting media**

Nutrient composition							
Media	Total Micro nutrients %			Total macro nutrients (ppm)			
	N	P	K	Zinc	Iron	Copper	Manganese
Vermicompost	0.640	0.61	0.54	68	3800	37.8	1000
Coir pith	0.224	0.21	0.62	32	420	13.6	200
Sugarcane node	0.210	0.22	0.86	58	104	11.8	57
Organic enricher	0.440	1.15	2.48	262	2200	46.7	664

Concrete differences in seedling vigour index and root length of sugarcane seedlings in different treatments were observed and presented in table 2 and plate1.

**Table2: Effect of rooting media on seedling vigour of sugarcane (Pooled data of 2014-15 and 2015-16)**

Treatment No.	Treatments	Root length (cm)	Seedling vigour index (SVI)
T1	Coir pith + Vermi compost (1:1)	17.25	11.86
T2	T1 + DAP (5kg / 100 kg media)	11.65	14.00
T3	<b>T1 + Urea (5kg / 100 kg media)</b>	<b>17.45</b>	<b>37.95</b>
T4	T1 + DAP (2.5 kg / 100 kg media) + Urea (2.5 kg / 100 kg media)	14.25	23.77
T5	T1 + Sett treatment with CaCl <sub>2</sub> (2g/lt) + Ethrel (2ml/lt)	15.05	19.61
T6	T1 + ZnSo <sub>4</sub> (0.5 kg) + FeSo <sub>4</sub> (1.0 kg) / 100 kg media)	15.98	19.01
T7	T1 + Sett treatment CaCl <sub>2</sub> (2g/lt) + Urea (10g/lt)	12.70	15.72
T8	T1 + Ethephon (2ml/lt)	14.15	26.94
T9	<b>T1 + Sett treatment with 10 % lime solution</b>	<b>18.30</b>	<b>33.36</b>
T10	<b>T1 + Sett treatment with 100 ppm GA3</b>	<b>15.86</b>	<b>30.46</b>
T11	T1 + Sett treatment with IAA 50 ppm	15.31	22.13
T12	T1 + Sett treatment with DM 45 + Chlorpyriphos	14.75	28.93
T13	Progressive farmers rooting media	17.80	24.84
	SEm+/-	1.63	2.20
	CD(0.05)	4.81	6.89



**Plate 1 – Photographs showing effect of rooting media on seedling vigour of sugarcane**

**Root length:** The root length of sugarcane seedling was more with the treatments T<sub>1</sub>(Coir pith + Vermi compost (1:1)), T<sub>3</sub> (T<sub>1</sub> + Urea (5kg / 100 kg media) , T<sub>9</sub> (T<sub>1</sub> + Sett treatment with 10 % lime solution) and T<sub>13</sub> (Progressive farmers rooting media) and these are statistical on par with each other.

**Seedling vigour index (SVI):** Seedling vigour index was high in the treatments T<sub>3</sub> (T<sub>1</sub> + Urea (5kg / 100 kg

media) , T<sub>9</sub> (T<sub>1</sub> + Sett treatment with 10 % lime solution) and T<sub>10</sub> (T<sub>1</sub> + Sett treatment with 100 ppm GA<sub>3</sub>) followed by T<sub>12</sub> (T<sub>1</sub> + Sett treatment with DM 45 + Chlorpyriphos), T<sub>8</sub>(T<sub>1</sub> + Ethephon (2ml/lt)) and T<sub>13</sub> (Progressive farmers rooting media). With the data we can derive that the treatments T<sub>3</sub> (T<sub>1</sub> + Urea (5kg / 100 kg media), T<sub>9</sub> (T<sub>1</sub> + Sett treatment with 10 % lime solution) and T<sub>10</sub> (T<sub>1</sub> + Sett treatment with 100 ppm

GA<sub>3</sub>) were found to perform better for sustainable sugarcane seedling vigour which can perform well for better establishment of sugarcane seedlings under transplanted fields. The treatments comprises of nitrogen in Urea, Calcium in 10% lime dipping, and growth substance GA<sub>3</sub> in addition to coir pith and vermin compost favored the germination per cent and improved the seedling vigour index in respective treatments over other treatments tested. Similar type

of observations with different rooting media was also reported with sugarcane seedling study by Vijakumar and Suresh, 2010 and Jain and Solomon, 2010 b.

**Acknowledgements:** The authors are sincerely thankful to Acharya N.G.Ranga Agricultural University, Guntur for giving permission and providing facilities to conduct the experiment at Regional Agricultural Research Station, Anakapalle during 2014-15 and 2015-16 cropping season.

### References:

- Jain R., S.Solomon, A.K.Srivastava and A.Chandra (2010 a). Sugarcane bud chips- A promising seed material. Sugar tech 12:67-69
- Mamta Prajapati, Smita Verma, Assessment of Mine Effluents on Water Quality ; Life Sciences international Research Journal , ISSN 2347-8691, Volume 2 Issue 1 (2015), Pg 329-331
- Jain R and S.Solomon (2010 b). Growth stimulating effect of ethephen on sprouting and early vigour of sugarcane bud chip. Sugar cane international 28 (3): 14-18
- Dr.Reddy, P.B , Sushma, P., The Impact of Life Style Factors on infertility And Reproductive Performance; Life Sciences International Research Journal , ISSN 2347-8691, Volume 2 Spl Issue (2015): Pg 37-44
- Naidu N.V., Sabitha N, and Raja Rajeswari.V (2015). Physiological and molecular approaches for enhancing sugarcane and sugar productivity Proceedings of 26<sup>th</sup> R&D sugarcane workers meet of A.P.Pg 76-80
- Dr.Chithambaram Chandrasekaran, Dr.D.Krupa Daniel, Craniosynostosis Study in South -East China; Life Sciences international Research Journal , ISSN 2347-8691, Volume 2 Issue 1 (2015), Pg 332-335
- Narsimha Rao G (1977). Chip bud method. The Hindu, July9, 1977
- Immanuel Rebecca, Patil Rohini Pradeep, Back to Kitchen: Food-Grade Media Is A Low-Cost Alternative to Commercial Bacteriological Media; Life Sciences International Research Journal , ISSN 2347-8691, Volume 2 Spl Issue (2015): Pg 49-55
- Prasada Rao. K (2015). Low cost technologies for productivity enchancement in sugarcane. Proceedings of 26<sup>th</sup> R&D sugarcane workers meet of A.P.Pg 70-75
- P.B.Reddy ,Jagdish Kannoja, Impact of Water Pollution on the Properties; Life Sciences International Research Journal , ISSN 2347-8691, Volume 1 Issue 1 (2014): Pg 47-50
- Ramaiah V.B., Narsimha Rao G, Prasad G.H(1977). Elimination of internode in sugarcane seed piece. Proceeding of international society of sugarcane technology: 1509-1513
- Rakesh Parmar , P.B.Reddy, Assessment of Surface Water Quality of Chambal; Life Sciences International Research Journal , ISSN 2347-8691, Volume 1 Issue 1 (2014): Pg 51-55
- Solomon S (2015). Revisiting cost effective technologies for sustainable sugarcane production. Proceedings of 26<sup>th</sup> R&D sugarcane workers meet of A.P.Pg 21-28
- Subhashree Dash, Bibhu Santosh Behera, Soubhagya Behera, Effective Management of Collar Rot of Groundnut Through Seedtreatment in Odisha; Life Sciences International Research Journal , ISSN 2347-8691, Volume 2 Issue 2 (2015): Pg 30-32
- Vijay Kumar M. and K.Suresh.(2010) Observational study on the identification of suitable media for raising seedlings with bud chips. Proceeding of 24<sup>th</sup> R&D sugarcane workers meet of A.P. Pg.110.
- Hanchinamani, C. N, Imamsaheb, S. J, Effect of Different Levels of Fertilizers and Spacing on Growth, Yield, Quality and Economics on onion (*Allium Cepa. L*) C V., Arkakalyan; Life Sciences International Research Journal , ISSN 2347-8691, Volume 2 Spl Issue (2015): Pg 46-51

Ch. Mukunda Rao, M. Bharathalakshmi, K. Veerabhadra Rao & N. Venugopala Rao  
Acharya N.G. Ranga Agricultural University, Regional Agricultural Research Station, Anakapalle – 531001, A.P.