

EFFECT OF CONCENTRATE SUPPLEMENTATION ON BROWSING BEHAVIOUR OF BERARI KIDS

Feeding behaviour in Berari kids

B. S. SADAR, D.R. AMBULKAR, T. D. JAISWAL

Abstract: Present experiment was carried out with the objective to study the effect of concentrate supplementation on browsing behaviour of Berari kids. Eighteen Berari kids (4 to 6 month of age) were allowed for browsing on natural vegetation around the farm premises from 10.00 -16.00 hours (6 hours daily). The kids were randomly distributed into three groups of equal number viz., T₀, T₁ and T₂. Kids of Gr-T₀ were allowed to browse on natural vegetation only. Kids of Gr-T₁ were supplemented grain mixture (16% crude protein and 65% total digestible nutrients) along with browsing whereas, Gr-T₂ were supplemented commercial concentrate feed (16% crude protein and 65% total digestible nutrients) in addition to browsing. The behaviour of animals was recorded once a week for 24 hours by direct visual observation. The kids of Gr-T₀ showed significantly (P<0.01) higher time on browsing activity (28.02 min/day) as compared to Gr-T₁ (23.38 min/day) and Gr-T₂ (22.69 min/day). A significant difference (P<0.01) was found in browsing time and ranged from 218.40 min/day in Gr-T₂ to 231.58 min/day in Gr-T₀. The ruminating time varied significantly (P<0.01) from 301.16 min/day in Gr-T₂ to 332.90 min/day in Gr-T₀. It was concluded that the concentrate supplementation reduced the grazing time, browsing time and ruminating time in kids.

Keywords: Behavior, Berari, Concentrate, Supplementation

Introduction: Major behaviours of goats when let loose for browsing in pasture are browsing, grazing, standing, idling, walking, ruminating, interaction and other drinking, scratching, self-grooming etc. The differences exist between in these behavioral activities across the breeds and within breed. These factors influence the amount of food consumed and therefore the net amount of nutrients available for metabolic ration. The present study was undertaken to record the behavior of recently registered Berari kids while grazing and in shelter.

Materials and Methods: The present experiment was conducted at "Berari Goat and Deccani Sheep Research, Demonstration and Training Centre, Borgaon Manju, Akola (Maharashtra). Eighteen Berari kids of 4 to 6 months age were selected irrespective of sex and randomly distributed into three equal groups. The kids were housed under loose housing system and let loose for browsing on natural vegetation around the farm premises for six hours (10:00 am - 4:00 pm). Kids of Gr T₀ were allowed to graze / browse on natural vegetation only. T₁ group was fed grain mixture (16% crude protein and 65% total digestible nutrients) along with grazing/browsing, whereas T₂ group was supplemented commercial concentrate (16% crude protein and 65% total digestible nutrients) in addition to grazing/browsing. Drinking water was provided *ad libitum* during the confined hours of the day to all the kids. The experiment was conducted for a period of two months during April and May (summer).

Behavioural recording (on 24 hr basis) was done once a week by direct visual observation. The activities like

grazing, browsing, ruminating and grazing: browsing ratios were recorded in terms of total time spent by individual kid (min/day). The data were analyzed with factorial completely randomized design (Snedecor and Cochran, 1994).

Results and Discussion: The meteorological observations during experimental period were presented in Table 1. The kids of Gr T₀ spent significantly (P<0.01) higher grazing and browsing time as compared to groups supplemented concentrate mixture (Table 2). It might be due to partial fulfillment of the dry matter requirement in groups (T₁ and T₂) by the concentrates. The grazing time showed decreasing trend during the progression of experiment. The increased environmental temperature might have been responsible for the gradual decrease in grazing time. Bordi et al. (1991) also reported higher grazing activity in control group (without any supplementation) than supplemented animals. The average browsing time was also lower in supplemented groups compared to control group. There was weekly significant decrease in average browsing time in all groups. On an average Berari kid spent 0.5% time on grazing and 35% time on browsing. Sharma et al. (1998) reported that Jamunapari and Barbari goats spent more time (52.9-62.0%) on browse during summer. Animut et al. (2005) found that time spent on browsing was 52.7% of active feeding time in sheep and goats when grazed together at three different stocking rates. Njoka-Njiru et al. (2001) also concluded that in the behaviour like feeding in dry period per 12 hrs observation time was 7.26 hrs in Toggenberg goats in subhumid Kenya.

Table 1. Meteorological observation during experimental period

Weeks	Ambient temperature (°C)		Sunshine hours
	Minimum	Maximum	
1	25.2	40.9	12.00
2	27.3	41.4	12.00

3	28.1	42.5	12.15
4	28.1	43.2	12.20
5	30.2	43.5	12.40
6	30.8	44.0	13.00
7	31.0	44.7	13.00
8	30.6	44.8	13.00

Table 2. Mean (±SE) grazing and browsing time of Berari goat kids fed during 10:00 am – 4:00 pm daily

Week	Grazing time (min/day)				Browsing time (min/day)			
	T ₀	T ₁	T ₂	Overall	T ₀	T ₁	T ₂	Overall
1	33.81±1.53	30.87±0.46	30.02±0.78	31.57±0.92 ^d	256.86±1.32	242.72±2.07	244.80±4.11	248.13±2.50 ^e
2	30.58±1.07	27.30±0.62	28.83±1.22	28.90±0.97 ^c	239.22±3.71	230.28±2.58	226.12±1.56	231.87±2.61 ^d
3	29.79±2.03	26.12±1.20	26.89±0.96	27.60±1.39 ^c	239.49±4.68	229.80±2.56	230.12±1.15	233.14±2.69 ^d
4	28.48±1.53	22.96±1.28	22.43± 1.01	24.62±1.27 ^b	231.38±5.29	221.86±4.05	219.86±2.91	224.36±4.08 ^c
5	26.95±1.49	21.89±1.49	19.97±1.24	22.94±1.40 ^b	228.27±4.95	211.62±3.01	210.35±1.18	216.74±3.04 ^b
6	25.86±2.13	19.26±1.58	18.44±1.00	21.18±1.53 ^a	224.95±3.82	207.49±4.05	210.47±1.72	214.30±3.19 ^b
7	23.65±1.98	19.06±1.21	16.96±0.52	19.89±1.23 ^a	216.65±2.76	208.40±2.39	206.52±2.22	210.52±2.45 ^a
8	25.07±1.68	19.62±0.80	17.98±0.58	20.89±3.06 ^a	215.79±2.26	205.24±0.87	198.99±2.06	206.67±1.73 ^a
Overall	28.02±1.17 ^b	23.38±1.53 ^a	22.69±1.84 ^a	24.70±1.51	231.58±4.79 ^b	219.68±4.81 ^a	218.40±5.24 ^a	223.22 ±4.95

Means bearing similar superscript does not differ significantly (P<0.01)

The overall grazing: browsing ratio for Berari kids when let loose for grazing during 10:00 am – 4:00 pm was 1: 8.29 for T₀, 1: 9.47 for T₁ and 1: 9.73 for T₂. As presented in Table 3, kids spent more time on browsing in supplemented groups (T₁ and T₂). Raats

(1988) found that with abundant forage, time spent browsing during the mornings is more than four times that of grazing (1: 4.2) and when forage is limited it decrease to equal time spent on browsing and grazing in South Africa.

Table 3. Total feeding (grazing + Browsing) time (min/day) of kids under different groups

Group	Week							
	1	2	3	4	5	6	7	8
T ₀	290.68	269.80	269.29	259.86	255.23	250.81	240.31	240.87
T ₁	273.59	257.58	255.92	244.83	233.51	226.75	227.46	224.87
T ₂	274.83	254.95	257.01	242.29	230.33	228.91	223.49	216.98

The overall average rumination time of Berari kids (Table 4) under grazing only group (T₀) was higher compared to the supplemented groups (T₁ and T₂). Forage intake has significant effect over the rumination time. As forage intake increased, there was significant increase in rumination time and when concentrate intake increased, there was reduction in rumination time (Shinde et al., 1997). The kids under grazing only group (T₀) spent more time for grazing

and browsing compared to supplemented groups (T₁ and T₂), it might lead to more consumption of forage than the other groups (T₁ and T₂) and therefore the time spent by kids in control group for rumination was more. Moon et al. (1994) studied the rumination behaviour over a 24 hr period with supplementary feeding of concentrate at various level and concluded that rumination time in was also significantly (P<0.01) affected by concentrate supplementation.

Table 4. Average Ruminating time (min/ day) of Berari goat kids

Week	T ₀	T ₁	T ₂	Overall
1	357.88±3.82 ^f	344.02±2.70 ^e	352.65±3.64 ^e	351.52±3.30 ^e
2	348.28±3.54 ^e	327.91±3.77 ^d	330.40±3.83 ^d	335.53±3.71 ^d
3	347.52±5.60 ^e	312.52±3.28 ^c	300.83±4.40 ^b	320.29±4.42 ^c
4	327.86±3.65 ^d	294.49±3.74 ^b	291.06±5.91 ^a	304.47±4.43 ^b
5	319.93±2.06 ^c	293.91±4.12 ^b	288.00±2.72 ^a	300.61±2.96 ^a
6	321.18±2.99 ^c	290.89±2.20 ^a	282.96±2.43 ^a	298.34±2.54 ^a
7	316.42±2.88 ^c	289.78±3.92 ^a	285.53±3.81 ^a	297.24±3.53 ^a
8	324.16±2.02 ^d	283.59±3.64 ^a	281.49±2.86 ^a	296.41±2.84 ^a
Overall	332.90±5.59 ^b	304.64±7.58 ^a	301.16±9.20 ^a	313.05 ±7.46

Means bearing similar superscript does not differ significantly (P<0.01)

It was concluded that the grazing time, browsing time and ruminating time were reduced in

concentrate supplemented kids. This might be because of adverse effect of environmental condition. Authors are thankful to

Acknowledgements: Associate Dean, Post graduate institute of veterinary and animal sciences, Akola for

financial support and to Deputy Director (Research), Berari Goat and Deccani Sheep Research, Demonstration and Training Centre, Borgaon Manju Dist. Akola for granting permission and facility for research work.

References:

1. Animut, G., Goetsch, A.L., Aiken, G.E., Puchala, R., Detweiler, C.R., Merkel, R.C., Sahl, T., Dawson, L.J., Johnson, Z.B. and Gipson, T.A. 2005. Grazing behavior and expenditure by sheep and goats co-grazing grass/forb pastures at three stocking rates. *Small Ruminant Research* 59: 191-201.
2. V.Gouri, T. Chitkaladevi, M. Bharatalakshmi, Balanced Fertilization For Higher Cane Yield in Sugarcane in North Coastal Zone of Andhra Pradesh; *Life Sciences International Research Journal* , ISSN 2347-8691, Volume 2 Spl Issue (2015): Pg 36-39
3. Bordi, A., Rosa, De G., Naoolitano, F., Vesce, G. and Rubino, R. 1994. Influence of feeding and supplementation on goats grazing behavior. *Options Mediterraneenes* 50: 57-68.
4. Moon, S., Enishi, O. and Hirota, H. 1994. Effect of supplementary concentrate on eating and rumination behavior in goats fed rye (*Secale cereal* l.) silage. *Nihon Chikusan Gakkaino* 65: 532-537.
5. Njoka-Njiru, E.N., Ojango, J.M., Ambula, M.K. and Ndirangu, C.M. 2001. Grazing behaviour of Sannen and Toggenburg goats in sub-humid conditions of Kenya. *Asian-Australian Journal of Animal Science* 14: 951-955.
6. Bikash Subba, S.K Ghosh, Population Dynamics of Lady Bird Beetle and Spiders in Relation to Weather Factors in tomato (*Lycopersicon Esculentum* L.) ; *Life Sciences International Research Journal* , ISSN 2347-8691, Volume 3 Issue 1 (2016): Pg 35-37
7. Raats, J.G. 1988. Feeding behaviour of free range goats. *South African Journal of Animal Science*. 18: 97-100.
8. Sharma, K., Saini, A.L., Singh, N. and Ogra, J.L. 1998. Seasonal variations in grazing behaviour and forage nutrient utilization by goats on a semi-arid reconstituted silvipasture. *Small Ruminant Research* 27: 47-54.
9. Shinde, A.K., Karim, S.A., Patnayak, B.C. and Mann, J.S. 1997. Dietary preference and grazing behaviour of sheep on *Cenchrus ciliaris* pasture in a semi-arid region of India. *Small Ruminant Research* 26: 119-122.
10. Sajad-Ul- Akbar Wani, Shehla Ishaque, Khurshed Ahmad, Status And Conservation of Waterfowl in Chatlam-Fushkooori Wetland Conservation Reserve, Pampore, Kashmir (Jammu & Kashmir) ; *Life Sciences International Research Journal* , ISSN 2347-8691, Volume 2 Spl Issue (2015): Pg 49-53
11. Snedecor, G.W. and Cochran, W.G. 1994. *Statistical Methods*. 8th ed., Affiliated East-West Press, New Delhi and Iowa State University Press, Iowa, USA.

B. S. Sadar, D.R. Ambulkar, T. D. Jaiswal

Department of Livestock Production & Management, Post Graduate Institute of Veterinary and Animal Sciences, Akola- 444 001 Maharashtra.