

## INFLUENCE OF DIFFERENT SUBSTRATES ON THE GROWTH AND YIELD OF OYSTER MUSHROOM

REEMA CHOURASIYA, PREETI MOYAL, KANUSHRI RANAWAT,  
RUPALI KAITHWAS, NEHA JOHRAM, REDDY P.B.

**Abstract:** The present investigation was carried out in Department of Microbiology at Government PG College, Ratlam, M.P (India). The farming of white rot edible fungus cultivation of *Pleurotus sajor-caju* was carried out on paddy straw, wheat straw and sugarcane bagasse using cylindrical block system to find out suitable substrate. The substrates significantly influenced on mycelial growth, spawn running, pin head formation, maturity of fruiting bodies, number of flushes and amount of yield. The highest number of primordia and fruiting bodies, and the amount of fresh weight was obtained with sugarcane bagasse in all flushes whereas, the lowest with wheat straw.

**Keywords:** Oyster mushroom, *Pleurotus sajor-caju*, Substrates, Yield.

**Introduction:** Oyster mushrooms are packed with most strong natural medicines for HIV and antioxidants (<https://www.awesomechef.in>). More than 100 varieties of species of mushrooms are being investigated for their health promoting benefits (Craig, W.J., 1999). of those hundred, about a half dozen really stand out for their ability to deliver a tremendous boost to your immune system (Borchers, A.T2004). Very recently, Barros, A.B., et al (2016) found a wide variety of health benefits of mushrooms like weight management, increasing vitamin D levels through your diet, improved nutrition, improved immune system function.

*Pleurotus* is renowned as an excellent mushroom. It can be cultured within a wide range of temperatures on different natural resources and agricultural wastes. Consequently, the cultivation of *P. sajor-caju* has increased extremely all over the world (Go, S.J et al, 1985, Pokhrel, C.P et al, 2013, Tupatkar, P.N. and Jadhao, S.M., 2006, Jhadav et al, 2014). Singh (1981) found that *P. sajor-caju* could be grown successfully on paddy straw at a temperature range of 19.1-30.50 C with 65-85% relative humidity. Whereas, Sivaprakasam and Kandaswamy (1981) reported a wide variety of substrates like waste papers, sugarcane bagasse, hulled maize cobs and rice straw for the production of its sporophores. In the present investigation, a white-rot fungal species *P. sajor-caju* were assessed for growth and yield on paddy straw, wheat straw and sugarcane bagasse.

**Substrate;** Choosing a correct substrate for mushroom cultivation is an important decision. A substrate is just any substance on which mycelium will grow. Carbon, nitrogen and inorganic compounds are required for proper growth of mushroom and cellulose, hemicelluloses and lignin form best source of carbon.

Large quantities of agriculture remains like rice, wheat and sugar cane are produced in India. Agriculture remains serve as important source of

lignocellulosic materials, which is best substrate for solid state fermentation of edible fungi such as *P. sajor-caju*.

At present, agriculture residues are disposed off through open-field burning, which causes a serious environment pollution problem. If such agricultural straw can support the growth of oyster mushroom, then it would be one of the solutions to convert these unpalatable wastes into a usual edible biomass of high market value, and serve as a cheap source of substrate for mushroom grower (Yang, W et al, 2013).

**Paddy straw:** The chemical composition of paddy straw varies between varieties and growing seasons, with higher nitrogen and cellulose contents in early-season rice compared to others. It contains 41% cellulose, 36% lignin, 0.45% total N. Wheat straw contains 1% protein, 13% lignin, 39% hemicelluloses, 40% cellulose (Sarnklong, C., et al, 2010).

**Wheat straw:** Wheat waste contains widely variable amounts of total nitrogen from 0.77-1.45%.

**Sugarcane bagasse:** Sugarcane bagasse contains 30% cellulose, 56% hemicelluloses, 13% lignin, 0.55% sugar, 36% Carbon, 0.448% nitrogen. The present investigation therefore aimed to study the influence of various agricultural wastes on the growth and yield of *P. sajor-caju* without any nutrient supplementation.

**Materials and Methods:** The required culture of oyster mushroom was obtained from Akshar Foods & Feeds Limited, Rasalpura, and Indore. (M.P.) and was grown on the potato dextrose agar (PDA 200 g/l diced potatoes; 20 g/l glucose, 15 g/l agar) medium at 25 °C for regular subculture. *P. sajor-caju* spawn was prepared in 850-ml polypropylene plastic bottles filled with cotton seed hull 87%, wheat bran 10%, sucrose 1%, plaster stone 1%, and calcium superphosphate 1% (w/w, in terms of dry weight) and subsequently sterilized at 121 °C for 80 min in autoclave. After cooling down to room temperature, the sterilized cotton seed hulls of every bottle were

inoculated with 5 cm<sup>2</sup> mycelial agar discs. The spawn was incubated in the laboratory at 26 ± 2 °C and 70% relative humidity for two weeks.

**Substrate preparation, inoculation and incubation:** The wheat straw, paddy straw, sugarcane bagasse, straws were completely dried under the Sun, and then chopped into small pieces of 1-2cm and soaked in water for overnight. After removal of the excess water, they were used as substrate in order to determine suitable substrates and suitable ratios for the farming of oyster mushroom. Dry weight of the substrates were noted before soaking and the bags full of different substrates weighed and were maintained at 1.5 Kg in a bag for each substrate. These inoculated bags were placed in growth room with 70-80% humidity and between 25-30°C temperatures, for impregnation of the substrate with mushroom spawn i.e. spawn running. Growth of mushroom was recorded daily for all the treatments. When such bags become full of growth and pin-heads started appearing, the bags were opened to facilitate the development of fruiting bodies. As soon as the fruiting bodies developed and attained their full size, they were cut just above surface of the substrate with sharp knife or blade. The yield parameters recorded were, days taken for the completion of spawn running, time (days) taken for the first appearance of pin-head formation, time (days) taken for the maturity of fruiting bodies, number of flushes, time interval (days) between flushes and yield of fresh mushroom on different crop wastes.

**Results And Discussion:** The summary of results regarding the completion of mycelium growth, spawn running, pin-head formation, maturity of fruiting bodies, number of flushes of fruiting bodies, and yield performance of *P. sajor-caju* on different substrates are mentioned in Table II. On sugarcane bagasse spawn running was completed earlier in 20 days followed by paddy straw. Wheat straw took maximum time period which was 43 days. In case of

pin-head formation sugarcane bagasse was better substrate (Table II). It was observed that time taken for first appearance of pinhead after spawning on the substrate was 23 days. It was followed by cotton waste 32 days. Maximum time 46 days were taken in case of wheat straw. In case of the maturity of fruiting bodies it was revealed that the minimum numbers of days were taken on the sugarcane bagasse (27 days), which proved to be the best substrate followed by rice straw (37.7 days). Maximum time period (50.7 days) was required for the maturity of fruiting bodies in case of wheat straw. Flushes of mushroom production varied among different substrates. The maximum flushes were on wheat and cotton waste (3.3). In case of *P.sajor-caju*, yield was 42.3 g on sugarcane bagasse.

Among the three substrates for the cultivation of oyster mushroom, sugar cane and rice straw gave the faster mycelial growth rate, time to primordial formation, time to first crop than wheat straw. In the present study oyster mushroom on rice straw and wheat straw substrate have less yield. Similar type of results were also obtained by many workers from different parts of the globe (Yang, W., et al, 2013, Naraian, R., et al, 2009, Zhang, R.Y., et al, 2014) but lower than some reported cultivation which have relatively good yields (Obodai and Vowotor, 2003). The reason may be physical nature and high C/N ratio of rice straw and wheat straw that were not suitable for the cultivars of oyster mushroom (Yang, W., et al, 2013, Girmay, Z., et al, 2016). Rice or wheat straw substrate in the present study was also found to be very vulnerable to drying with high porosity as reported by Wang, L. et al, (2016) which affected sporophore formation.

Based on the investigations, it can be concluded that yield of oyster mushroom, *P. sajor-caju* appeared earlier on sugarcane bagasse followed by rice straw and the maximum flushes were obtained from wheat straw. The minimum flush was taken on paddy straw and the maximum yield on fresh basis was obtained from wheat straw.

## References:

1. Craig, W.J., 1999. Health-promoting properties of common herbs. The American journal of clinical nutrition, 70(3), pp.491s-499s).
2. Borchers, A.T., Keen, C.L. and Gershwin, M.E., 2004. Mushrooms, tumors, and immunity: an update. Experimental Biology and Medicine, 229(5), pp.393-406.
3. Barros, A.B., Bell, V., Ferrão, J., Calabrese, V. and Fernandes, T.H., 2016. Mushroom Biomass: Some Clinical Implications of  $\beta$ -Glucans and Enzymes. Current Research in Nutrition and Food Science Journal, 4(Special Issue Nutrition in Conference October 2016), pp.37-47.
4. Pokhrel, C.P., Kalyan, N., Budathoki, U. and Yadav, R.K.P., 2013. Cultivation of *Pleurotus sajor-caju* using different agricultural residues. International Journal of Agricultural Policy and Research, 1(2), pp.19-23.
5. Tupatkar, P.N. and Jadhao, S.M., 2006. Effects of different substrates on yield of oyster mushroom (*Pleurotus sajor-caju*). Agric Sci Digest, 26(3), pp.224-226.
6. Jadhav, Ashok, C., Dattatray, B. Shinde, Santosh, B. Nadre and Darshan, S. Deore. 2014. Quality improvement of casing material and yield in milky mushroom (*Calocybe indica*) by using

- biofertilizers and different substrates. In Proceedings of 8th International Conference on Mushroom Biology and Mushroom Products (ICMBMP8), New Delhi, India, 19-22 November 2014. Volume I & II, pp. 359-364. ICAR-Directorate of Mushroom Research, 2014.
7. Go, S.J., Shin, G.C. and Yoo, Y.B., 1985. Protoplast formation, regeneration and reversion in *Pleurotus ostreatus* and *P. sajor-caju*. The Korean Journal of Mycology, 13(3), pp.169-177.
  8. Dr.Chithambaram Chandrasekaran, Dr.D.Krupa Daniel, *Craniosynostosis Study in South -East China; Life Sciences international Research Journal* , ISSN 2347-8691, Volume 2 Issue 1 (2015), Pg 332-335
  9. Sarnklong, C., Cone, J.W., Pellikaan, W. and Hendriks, W.H., 2010. Utilization of rice straw and different treatments to improve its feed value for ruminants: a review. Asian-Australasian Journal of Animal Sciences, 23(5), p.680.
  10. Yang, W., Guo, F. and Wan, Z., 2013. Yield and size of oyster mushroom grown on rice/wheat straw basal substrate supplemented with cotton seed hull. Saudi journal of biological sciences, 20(4), pp.333-338.
  11. Yang, W., Guo, F. and Wan, Z., 2013. Yield and size of oyster mushroom grown on rice/wheat straw basal substrate supplemented with cotton seed hull. Saudi journal of biological sciences, 20(4), pp.333-338.
  12. Naraian, R., Sahu, R.K., Kumar, S., Garg, S.K., Singh, C.S. and Kanaujia, R.S., 2009. Influence of different nitrogen rich supplements during cultivation of *Pleurotus florida* on corn cob substrate. The Environmentalist, 29(1), p.1.
  13. Zhang, R.Y., Hu, D.D., Ma, X.T., Li, S.G., Gu, J.G. and Hu, Q.X., 2014. Adopting stick spawn reduced the spawn running time and improved mushroom yield and biological efficiency of *Pleurotus eryngii*. Scientia Horticulturae, 175, pp.156-159.
  14. Reddy P.B, G.R.Gangle, Evaluation of the Function and Impact of Hocky Stick Data of the intergovernmental Panel on Climate Change; Life Sciences International Research Journal , ISSN 2347-8691, Volume 2 Issue 2 (2015): Pg 133-135
  15. Obodai and Vowotor, M.C.O. Obodai, K.A. Vowotor, 2003. Comparative study on the growth and yield of *Pleurotus ostreatus* mushroom on different lignocellulosic by-products. J. Ind. Microb. Biotechnol. 30 (2003), pp. 146-149.
  16. Girmay, Z., Gorems, W., Birhanu, G. and Zewdie, S., 2016. Growth and yield performance of *Pleurotus ostreatus* (Jacq. Fr.) Kumm (oyster mushroom) on different substrates. AMB Express, 6(1), p.87.
  17. Wang, L., Mao, J., Zhao, H., Li, M., Wei, Q., Zhou, Y. and Shao, H., 2016. Comparison of characterization and microbial communities in rice straw-and wheat straw-based compost for *Agaricus bisporus* production. Journal of Industrial Microbiology & Biotechnology, 43(9), pp.1249-1260.
  18. Swapnil Lokegaonkar, Bela Nabar, Statistical Survey of Rural and Urban Population of Thane District for Putative Dental Plaque Risk Factors; Life Sciences International Research Journal , ISSN 2347-8691, Volume 2 Spl Issue (2015): Pg 211-218

Reema Chourasiya, Preeti Moyal, , Rupali Kaithwas, Neha Johram, Reddy P.B, Department of Microbiology, Kanushri Ranawat Department of Biotechnology, Govt.PG College, Ratlam.

**Table I: Chemical composition of Substrates**

Substrate	Chemical composition (% w/w)			
	Cellulose	Hemi cellulose	Lignin	Total nitrogen
Sugarcane bagasse	8.34	30	56	0.448
Wheat straw	40	39	13	0.77
Paddy straw	39.2	23.5	36.5	0.457

**Table II: Time taken for growth and yield of *P.sajor-caju* on different substrates**

Substrates	Spawn running (in days)	Pin head formation (in days)	Maturation of fruiting bodies (in days)	Number of flushes	Yield (g)
Wheat straw	44.2a	46.8a	51.1a	3.4a	38.6a
Paddy straw	41.5a	44.2a	49.8a	3.1a	40.4a
Sugarcane bagasse	20.0a	24.0a	26.8a	3.0a	41.1a